

PRODUCTS AND H-HOMOGENEITY

ANDREA MEDINI

ABSTRACT. Building on work of Terada, we prove that h-homogeneity is productive in the class of zero-dimensional spaces. Then, by generalizing a result of Motorov, we show that for every zero-dimensional space X there exists a zero-dimensional space Y such that $X \times Y$ is h-homogeneous. Also, we simultaneously generalize results of Motorov and Terada by showing that if X is a zero-dimensional space such that the isolated points are dense then X^κ is h-homogeneous for every infinite cardinal κ . Finally, we show that a question of Terada (whether X^ω is h-homogeneous for every zero-dimensional first-countable X) is equivalent to a question of Motorov (whether such an infinite power is always divisible by 2) and give some partial answers.

All spaces in this paper are assumed to be Tychonoff. For all undefined topological notions, see [5].

Definition 1. A space X is *h-homogeneous* (or *strongly homogeneous*) if all non-empty clopen subsets of X are homeomorphic to each other.

The Cantor set, the rationals and the irrationals are examples of h-homogeneous spaces. Every connected space is h-homogeneous. A finite space is h-homogeneous if and only if it has size at most 1. The concept of h-homogeneity has been studied (mostly in the zero-dimensional case) by several authors: see [9] for an extensive list of references.

We will denote by $\text{CO}(X)$ the Boolean algebra of the clopen subsets of X . If X is h-homogeneous then $\text{CO}(X)$ is homogeneous; the converse holds if X is compact and zero-dimensional (see [8]).

1. THE PRODUCTIVITY OF H-HOMOGENEITY

All spaces in this section are assumed to be infinite.

In [16], the productivity of h-homogeneity is stated as an open problem (see also [9] and [10]), and the following is proved (see theorem 3.3).

Theorem 2 (Terada). *If X_i is h-homogeneous and zero-dimensional for every $i \in I$ and $X = \prod_{i \in I} X_i$ is compact or non-pseudocompact, then X is h-homogeneous.*

We will show that the requirement that X is compact or non-pseudocompact can actually be dropped (see theorem 10) and that if X is pseudocompact then the zero-dimensionality requirement can be dropped (see theorem 9).

For proofs of the following basic facts about βX , see section 11 in [11]. Given any open subset U of X , define $\text{Ex}(U) = \beta X \setminus \text{cl}_{\beta X}(X \setminus U)$. It is easy to see that $\text{Ex}(U)$ is the biggest open subset of βX such that its intersection with X is U . If C is a clopen subset of X then $\text{Ex}(C) = \text{cl}_{\beta X}(C)$, hence $\text{Ex}(C)$ is clopen in βX . Furthermore, the collection $\{\text{Ex}(U) : U \text{ is open in } X\}$ is a base for βX .

Date: November 5th, 2009.

Remark. It is not true that βX is zero-dimensional whenever X is zero-dimensional (see example 6.2.20 in [5] or 3.39 in [17]). If βX is zero-dimensional then X is called *strongly zero-dimensional*.

Proposition 3. *Assume that $X \times Y$ is pseudocompact. If $C \subseteq X \times Y$ is clopen then C can be written as the union of finitely many open rectangles.*

Proof. By Glicksberg's theorem (see theorem 1 in [7] or theorem 8.25 in [17]), $X \times Y$ is C^* -embedded in $\beta X \times \beta Y$. By the universal property of the Stone-Čech compactification, there exists a homeomorphism $h : \beta X \times \beta Y \rightarrow \beta(X \times Y)$ such that $h(x, y) = (x, y)$ whenever $(x, y) \in X \times Y$.

Since $\{\text{Ex}(U) : U \text{ is open in } X\}$ is a base for βX and $\{\text{Ex}(V) : V \text{ is open in } Y\}$ is a base for βY , the collection

$$\mathcal{B} = \{\text{Ex}(U) \times \text{Ex}(V) : U \text{ is open in } X \text{ and } V \text{ is open in } Y\}$$

is a base for $\beta X \times \beta Y$. Therefore $\{h[B] : B \in \mathcal{B}\}$ is a base for $\beta(X \times Y)$, hence we can write $\text{Ex}(C) = h[B_1] \cup \dots \cup h[B_n]$ for some $B_1, \dots, B_n \in \mathcal{B}$ by compactness.

Finally, if we let $B_i = \text{Ex}(U_i) \times \text{Ex}(V_i)$ for each i , we get

$$\begin{aligned} C &= \text{Ex}(C) \cap X \times Y \\ &= (h[B_1] \cup \dots \cup h[B_n]) \cap h[X \times Y] \\ &= h[B_1 \cap X \times Y] \cup \dots \cup h[B_n \cap X \times Y] \\ &= (B_1 \cap X \times Y) \cup \dots \cup (B_n \cap X \times Y) \\ &= (U_1 \times V_1) \cup \dots \cup (U_n \times V_n), \end{aligned}$$

that concludes the proof. $\square$

Lemma 4. *Assume that $C \subseteq X \times Y$ is a clopen set that can be written as the union of finitely many rectangles. Then C can be written as the union of finitely many pairwise disjoint clopen rectangles.*

Proof. For every $x \in X$, let $C_x = \{y \in Y : (x, y) \in C\}$ be the corresponding vertical cross-section. For every $y \in Y$, let $C^y = \{x \in X : (x, y) \in C\}$ be the corresponding horizontal cross-section. Since C is clopen, each cross-section is clopen.

Let $\mathcal{A}$ be the Boolean subalgebra of $\text{CO}(X)$ generated by $\{C^y : y \in Y\}$. Since $\mathcal{A}$ is finite, it must be atomic. Let $P_1, \dots, P_m$ be the atoms of $\mathcal{A}$. Similarly, let $\mathcal{B}$ be the Boolean subalgebra of $\text{CO}(Y)$ generated by $\{C_x : x \in X\}$, and let $Q_1, \dots, Q_n$ be the atoms of $\mathcal{B}$.

Observe that the rectangles $P_i \times Q_j$ are clopen and pairwise disjoint. Furthermore, given any i, j , either $P_i \times Q_j \subseteq C$ or $P_i \times Q_j \cap C = \emptyset$. Hence C is the union of a (finite) collection of such rectangles. $\square$

Proposition 5. *Assume that $X \times Y$ is pseudocompact. If X is h -homogeneous and Y is h -homogeneous then $X \times Y$ is h -homogeneous.*

Proof. If X and Y are both connected then $X \times Y$ is connected. So assume without loss of generality that X is not connected. Since X is also h -homogeneous, it follows that $X \cong n \times X$ for every $n \in \omega \setminus \{0\}$. Therefore $X \times Y \cong n \times X \times Y$ for every $n \in \omega \setminus \{0\}$.

Now let $C \subseteq X \times Y$ be clopen and non-empty. By proposition 3 and lemma 4, we can write C as the disjoint union of finitely many, say n , non-empty clopen rectangles. By the h -homogeneity of X and Y , each such rectangle is homeomorphic to $X \times Y$. Therefore $C \cong n \times X \times Y \cong X \times Y$. $\square$

Corollary 6. *Assume that $X = X_1 \times \cdots \times X_n$ is pseudocompact. If each X_i is h -homogeneous then X is h -homogeneous.*

An obvious modification of the proof of proposition 3 yields the following.

Proposition 7. *Assume that $X = \prod_{i \in I} X_i$ is pseudocompact. If $C \subseteq X$ is clopen then C can be written as the union of finitely many open rectangles.*

Corollary 8. *Assume that $X = \prod_{i \in I} X_i$ is pseudocompact. If $C \subseteq X$ is clopen then C depends on finitely many coordinates.*

Theorem 9. *Assume that $X = \prod_{i \in I} X_i$ is pseudocompact. If X_i is h -homogeneous for every $i \in I$ then X is h -homogeneous.*

Proof. Let $C \subseteq X$ be clopen and non-empty. By corollary 8, there exists a finite subset F of I such that C is homeomorphic to $C' \times \prod_{i \in I \setminus F} X_i$, where C' is a clopen subset of $\prod_{i \in F} X_i$. Since $\prod_{i \in F} X_i$ is h -homogeneous by corollary 6, we must have $C' \cong \prod_{i \in F} X_i$. Hence $C \cong X$. $\square$

Theorem 10. *If X_i is h -homogeneous and zero-dimensional for every $i \in I$ then $X = \prod_{i \in I} X_i$ is h -homogeneous.*

Proof. If X is pseudocompact, apply theorem 9; if X is non-pseudocompact, apply theorem 2. $\square$

2. SOME APPLICATIONS

The following theorem was proved by Terada in [16] under the additional assumption that X is zero-dimensional (see theorem 2.4).

Theorem 11 (Terada). *Assume that X is non-pseudocompact and it has a π -base $\mathcal{B}$ consisting of clopen sets that are homeomorphic to X . Then X is h -homogeneous.*

Terada's proof uses the fact that a zero-dimensional non-pseudocompact space can be written as the disjoint union of infinitely many of its non-empty clopen subsets (the converse is also true, trivially). However, that is the only consequence of zero-dimensionality that is actually used. Therefore such assumption is redundant by the following lemma.

Lemma 12. *If X is non-pseudocompact and it has a π -base $\mathcal{B}$ consisting of clopen sets then X can be written as the disjoint union of infinitely many of its non-empty clopen subsets.*

Proof. Let $f : X \rightarrow \mathbb{R}$ be an unbounded continuous function. Then the range of f contains a countably infinite subset D that is closed and discrete as a subset of $\mathbb{R}$. Let $D = \{d_n : n \in \omega\}$ be an enumeration without repetitions and choose pairwise disjoint open sets U_n such that $d_n \in U_n$ for each $n \in \omega$. Then choose open sets V_n such that $\overline{V_n} \subseteq U_n$ and $\text{diam}(V_n) \leq 2^{-n}$ for each $n \in \omega$. Finally, let $B_n \in \mathcal{B}$ such that $B_n \subseteq f^{-1}[V_n]$ for each $n \in \omega$. It is easy to check that $B = \bigcup_{n \in \omega} B_n$ is clopen. Hence $X \setminus B, B_0, B_1, \dots$ is the desired partition of X . $\square$

The following result was proved by Motorov under the additional assumption that X is compact and first-countable (see theorem 2 in [12]), though the assumption of first-countability is not actually used in the proof.

Theorem 13. *Assume that Y has a π -base $\mathcal{B}$ consisting of clopen sets. Then $X = (Y \times 2 \times \prod \mathcal{B})^\kappa$ is h -homogeneous for every infinite cardinal κ .*

Proof. An analysis of Motorov's proof shows that the only consequence of the compactness of X that is used is the fact that clopen sets in X depend on finitely many coordinates. Therefore the same proof works if X is pseudocompact by corollary 8.

Now assume that X is non-pseudocompact. One can easily check that X has a π -base consisting of clopen sets that are homeomorphic to X . Therefore X is h -homogeneous by theorem 11. $\square$

Corollary 14. *For every zero-dimensional space X there exists a zero-dimensional space Y such that $X \times Y$ is h -homogeneous.*

The following is due to Matveev (see proposition 3 in [9]).

Proposition 15 (Matveev). *Assume that X is a zero-dimensional space with a π -base $\mathcal{B}$ consisting of clopen sets that are homeomorphic to X . If there exists a sequence $\langle U_n : n \in \omega \rangle$ of open sets in X that converges to a point then X is h -homogeneous.*

The case $\kappa = \omega$ of the following result is an easy consequence of proposition 15. Motorov first proved it under the additional assumption that X is first-countable and compact (see theorem 1 in [12]). Terada proved it for an arbitrary infinite κ , under the additional assumption that X is non-pseudocompact (see corollary 3.2 in [16]).

Theorem 16. *Assume that X is a zero-dimensional space such that the isolated points are dense in X . Then X^κ is h -homogeneous for every infinite cardinal κ .*

Proof. Since $X^\kappa = (X^\omega)^\kappa$ for every infinite cardinal κ , we can assume that $\kappa = \omega$ by theorem 10.

Let D be the set of isolated points of X and let $\text{Fn}(\omega, D)$ be the set of finite partial functions from ω to D . Given $s \in \text{Fn}(\omega, D)$, define $U_s = \{f \in X^\omega : f \supseteq s\}$. Now fix $d \in D$ and let $f \in X^\omega$ be the constant function with value d . It is easy to see that $\langle U_{f \upharpoonright n} : n \in \omega \rangle$ is a sequence of open sets in X^ω that converges to f . Furthermore $\mathcal{B} = \{U_s : s \in \text{Fn}(\omega, D)\}$ is a π -base for X^ω consisting of clopen sets that are homeomorphic to X^ω . So X^ω is h -homogeneous by proposition 15. $\square$

3. INFINITE POWERS OF ZERO-DIMENSIONAL FIRST-COUNTABLE SPACES

It is a well-known fact that every zero-dimensional first-countable h -homogeneous space is homogeneous. As announced by Motorov in [13], the converse holds for zero-dimensional first-countable compact spaces of uncountable cellularity (see [6] for a proof). In [1], van Douwen constructed a zero-dimensional first-countable compact homogeneous space that is not h -homogeneous (actually, it has no proper subspaces homeomorphic to itself). In [14], using similar techniques, Motorov constructed a zero-dimensional first-countable compact homogeneous space that is not divisible by 2 (in the sense of definition 18).

In [16], Terada asked whether X^ω is h -homogeneous for every zero-dimensional first-countable X . In [2], the following remarkable theorem is proved.

Theorem 17 (Dow and Pearl). *If X is a zero-dimensional first-countable space then X^ω is homogeneous.*

However, Terada's question remains open. In [12] and [13], Motorov asks whether such an infinite power is always divisible by 2. Using theorem 17, we will show that the two questions are equivalent: actually even weaker conditions suffice (see proposition 21).

Definition 18. A space F is a *factor* of X (or X is *divisible* by F) if there exists Y such that $F \times Y \cong X$. If $F \times X \cong X$ then F is a *strong factor* of X (or X is *strongly divisible* by F).

We will use the following lemma freely in the rest of this section. It shows that divisibility and strong divisibility are the same for infinite powers.

Lemma 19. *The following are equivalent.*

- (1) F is a factor of X^ω .
- (2) $F \times X^\omega \cong X^\omega$.
- (3) $F^\omega \times X^\omega \cong X^\omega$.

Proof. The implications $2 \rightarrow 1$ and $3 \rightarrow 1$ are clear.

Assume 1. Then there exists Y such that $F \times Y \cong X^\omega$, hence

$$X^\omega \cong (X^\omega)^\omega \cong (F \times Y)^\omega \cong F^\omega \times Y^\omega.$$

Since multiplication by F or by F^ω does not change the right hand side, it follows that 2 and 3 hold. $\square$

Lemma 20. *Assume that Y is a non-empty first-countable zero-dimensional space. Then $X = (Y \oplus 1)^\omega$ is h-homogeneous and $X \cong Y^\omega \times (Y \oplus 1)^\omega \cong 2^\omega \times Y^\omega$.*

Proof. Recall that $1 = \{0\}$ and let $g \in X$ be the constant function with value 0. For each $n \in \omega$, define

$$U_n = \{f \in X : f(i) = 0 \text{ for all } i < n\}.$$

Observe that $\mathcal{B} = \{U_n : n \in \omega\}$ is a local base for X at g consisting of clopen sets that are homeomorphic to X . But X is homogeneous by theorem 17, therefore it has a base (hence a π -base) consisting of clopen sets that are homeomorphic to X . It follows from proposition 15 that X is h-homogeneous.

To prove the second statement, observe that

$$X \cong (Y \oplus 1) \times X \cong (Y \times X) \oplus X,$$

hence $X \cong Y \times X$ by h-homogeneity. It follows that $X \cong Y^\omega \times (Y \oplus 1)^\omega$. Finally,

$$Y^\omega \times (Y \oplus 1)^\omega \cong (Y^\omega \times (Y \oplus 1))^\omega \cong (Y^\omega \oplus Y^\omega)^\omega \cong 2^\omega \times Y^\omega,$$

that concludes the proof. $\square$

Proposition 21. *Assume that X is a first-countable zero-dimensional space containing at least two points. Then the following are equivalent.*

- (1) $X^\omega \cong (X \oplus 1)^\omega$
- (2) $X^\omega \cong Y^\omega$ for some Y with at least one isolated point.
- (3) X^ω is h-homogeneous.
- (4) X^ω has a clopen subset that is strongly divisible by 2.
- (5) X^ω has a proper clopen subspace homeomorphic to X^ω .
- (6) X^ω has a proper clopen subspace as a factor.

Proof. The implication $1 \rightarrow 2$ is trivial; the implication $2 \rightarrow 3$ follows from lemma 20; the implications $3 \rightarrow 4 \rightarrow 5 \rightarrow 6$ are trivial.

Assume that 6 holds. Let C be a proper clopen subset of X^ω that is also a factor of X^ω and let $D = X^\omega \setminus C$. Then

$$\begin{aligned} X^\omega &\cong (C \oplus D) \times X^\omega \\ &\cong (C \times X^\omega) \oplus (D \times X^\omega) \\ &\cong X^\omega \oplus (D \times X^\omega) \\ &\cong (1 \oplus D) \times X^\omega, \end{aligned}$$

hence $X^\omega \cong (1 \oplus D)^\omega \times X^\omega$. Since $(1 \oplus D)^\omega \cong 2^\omega \times D^\omega$ by lemma 20, it follows that $X^\omega \cong 2^\omega \times X^\omega$. Therefore 1 holds by lemma 20. $\square$

The next two propositions show that in the pseudocompact case we can say something more.

Proposition 22. *Assume that X is a first-countable zero-dimensional space such that X^ω is pseudocompact. Then $C^\omega \cong (X \oplus 1)^\omega$ for every non-empty proper clopen subset C of X^ω .*

Proof. Let C be a non-empty proper clopen subset of X^ω . It follows from corollary 8 that $C \cong C \times X^\omega$, hence $C^\omega \cong C^\omega \times X^\omega$. Since $C^\omega \times X^\omega$ clearly has a proper clopen subset homeomorphic to itself, proposition 21 implies that C^ω is strongly divisible by 2. Since $2^\omega \times X^\omega \cong (X \oplus 1)^\omega$ by lemma 20, it follows that $C^\omega \cong C^\omega \times (X \oplus 1)^\omega$.

On the other hand, $(X \oplus 1)^\omega \cong X^\omega \times (X \oplus 1)^\omega$ by lemma 20. Hence $(X \oplus 1)^\omega$ has a clopen subset homeomorphic to $C \times (X \oplus 1)^\omega$. But lemma 20 shows that $(X \oplus 1)^\omega$ is h-homogeneous, so $C \times (X \oplus 1)^\omega \cong (X \oplus 1)^\omega$. Therefore $C^\omega \times (X \oplus 1)^\omega \cong (X \oplus 1)^\omega$, that concludes the proof. $\square$

Proposition 23. *In addition to the hypotheses of proposition 21, assume that X^ω is pseudocompact. Then the following can be added to the list of equivalent conditions.*

- (7) X^ω has a proper clopen subspace C such that $C \cong Y^\omega$ for some Y .

Proof. The implication $5 \rightarrow 7$ is trivial.

Assume that 7 holds. Then clearly $C^\omega \cong C$. Therefore $C \cong (X \oplus 1)^\omega$ by proposition 22. Hence C is strongly divisible by 2 by lemma 20, showing that 4 holds. $\square$

Finally, we point out that proposition 21 can be used to give a positive answer to Terada's question for a certain class of spaces. We will need the following definition (see section 1 in [3]).

Definition 24. A space X is *ultraparacompact* if every open cover of X has a refinement consisting of pairwise disjoint clopen sets.

A metric space X is ultraparacompact if and only if $\dim X = 0$ (see the corollary to theorem II.6 in [15]). For such a metric space X , van Engelen proved that X^ω is h-homogeneous if X is first-category in itself or X has a completely metrizable dense subset (see theorems 4.2 and 4.4 in [4]).

Proposition 25. *Assume that X is a zero-dimensional first-countable space. If X^ω is ultraparacompact and non-Lindelöf then X^ω is h-homogeneous.*

Proof. Let $\mathcal{U}$ be an open cover of X^ω with no countable subcovers. By ultraparacompactness, there exists a refinement $\mathcal{V}$ of $\mathcal{U}$ consisting of pairwise disjoint clopen sets. Let $\mathcal{V} = \{C_\alpha : \alpha \in \kappa\}$ be an enumeration without repetitions, where κ is an uncountable cardinal.

Now fix $x \in X^\omega$ and a local base $\{U_n : n \in \omega\}$ at x consisting of clopen sets. Since X^ω is homogeneous by theorem 17, for each $\alpha < \kappa$ we can find $n(\alpha) \in \omega$ such that an homeomorphic clopen copy $U'_{n(\alpha)}$ of $U_{n(\alpha)}$ is contained in C_α . Fix $S \subseteq \kappa$ of size κ such that $n(\alpha) = n(\beta)$ for every $\alpha, \beta \in S$. Clearly $\bigcup_{\alpha \in S} U'_{n(\alpha)}$ is a clopen subset of X^ω that is strongly divisible by 2. So X^ω is h-homogeneous by proposition 21. $\square$

Acknowledgement. The author thanks his advisor Ken Kunen for valuable comments on an earlier version of this paper.

REFERENCES

- [1] E. K. VAN DOUWEN. A compact space with a measure that knows which sets are homeomorphic. *Adv. in Math.* **52:1** (1984), 1–33.
- [2] A. DOW AND E. PEARL. Homogeneity in powers of zero-dimensional first-countable spaces. *Proc. Amer. Math. Soc.* **125** (1997), 2503–2510.
- [3] R. L. ELLIS. Extending continuous functions on zero-dimensional spaces. *Math. Ann.* **186** (1970), 114–122.
- [4] F. VAN ENGELEN. On the homogeneity of infinite products. *Topology Proc.* **17** (1992), 303–315.
- [5] R. ENGELKING. *General topology*. Sigma Series in Pure Mathematics, vol. 6. Heldermann Verlag, Berlin, 1989.
- [6] S. SHELAH AND S. GESCHKE. Some notes concerning the homogeneity of Boolean algebras and Boolean spaces. *Topology Appl.* **133:3** (2003), 241–253.
- [7] I. GLICKSBERG. Stone-Čech compactifications of products. *Trans. Amer. Math. Soc.* **90** (1959), 369–382.
- [8] S. KOPPELBERG. *Handbook of Boolean algebras. Volume 1*. Edited by J. D. Monk and R. Bonnet. North-Holland, Amsterdam, 1989.
- [9] M. V. MATVEEV. Basic homogeneity in the class of zero-dimensional spaces. *Filomat.* **12:2** (1998), 13–20.
- [10] M. V. MATVEEV AND S. A. SHKARIN. Some questions on topological homogeneity. *Questions and Answers in General Topology.* **18:1** (2000), 1–16.
- [11] J. VAN MILL. An invitation to $\beta\omega$. Available at <http://www.few.vu.nl/~vanmill/teaching/caput/caput.pdf>.
- [12] D. B. MOTOROV. Homogeneity and π -networks. *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* **4** (1989), 31–34 (in Russian); English translation in: *Moscow Univ. Math. Bull.* **44:4** (1989), 45–50.
- [13] D. B. MOTOROV. Zero-dimensional and linearly ordered bicomacta: properties of homogeneity type. *Uspekhi Math. Nauk.* **44:6** (1989), 159–160 (in Russian); English translation in: *Russ. Math. Surv.* **44:6** (1989), 190–191.
- [14] D. B. MOTOROV. A homogeneous zero-dimensional bicomactum which is not divisible by 2. *General topology. Spaces, mappings and functors. Collection of scientific works*. Edited by V. V. Fedorchuk et al. Moskva: Izdatel'stvo Moskovskogo Universiteta. (1992), 65–73 (in Russian).
- [15] J.-I. NAGATA. *Modern dimension theory*. Revised edition. Sigma Series in Pure Mathematics, vol. 2. Heldermann Verlag, Berlin, 1983.
- [16] T. TERADA. Spaces whose all nonempty clopen subsets are homeomorphic. *Yokohama Math. Jour.* **40** (1993), 87–93.
- [17] R. C. WALKER. *The Stone-Čech compactification*. Springer-Verlag, New York, 1974.

UNIVERSITY OF WISCONSIN-MADISON MATHEMATICS DEPT.
E-mail address: medini@math.wisc.edu